

External malformations in the green sea urchin *Arbacia dufresnii* (Blainville, 1825) in populations from Golfo Nuevo, Patagonia, Argentina

Florencia DI MARCO^{1,2*}, Sebastián GIACOMINO¹, Laura LÓPEZ GRECO^{2,3,4}
& Ximena GONZÁLEZ PISANI^{1,2}

¹Centro para el Estudio de Sistemas Marinos (CESIMAR), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Centro Nacional Patagónico (CENPAT), Blvd. Almte. Brown 2915 (U9120ACD), Puerto Madryn, Argentina. ²Laboratorio de Ecotoxicología en Invertebrados Acuáticos (LEIA), Instituto Patagónico del Mar (IPaM), Universidad Nacional de la Patagonia San Juan Bosco (UNPSJB), Bv. Almte. Brown 3051 (9120), Puerto Madryn, Argentina. ³Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Biodiversidad y Biología Experimental (DBBE FCEN-UBA), Laboratorio de Biología de la Reproducción, Crecimiento y Nutrición de Crustáceos Decápodos, Ciudad Universitaria, Buenos Aires, Argentina. ⁴CONICET - Universidad de Buenos Aires, Instituto de Biodiversidad y Biología Experimental y Aplicada (IBBEA), Buenos Aires, Argentina. *Corresponding author:* Lic. Florencia Di Marco, Bvd. Brown 2915-U9120ACD - Puerto Madryn – Chubut, phone (+54) 0280 488-3184 (Int 1301) and fax number (+54) 0280 488-3543, e-mail: flordm6@gmail.com / fdimarco@cenpat-conicet.gob.ar

Abstract: Malformations in sea urchins can manifest in various forms, often influenced by environmental and biological factors. Adult *Arbacia dufresnii* (Blainville, 1825) exhibit pentaradial symmetry, spines in the ambulacral areas, absence of spines in the interambulacral areas, and a rhomboidal periproct composed of four ossicles. This report aims to document the presence of external malformations in the green sea urchin *Arbacia dufresnii* in the population of Golfo Nuevo, Patagonia, Argentina (42°42' S 64°30' W). The specimens were collected at Golfo Nuevo. A total of 319 specimens were collected, out of which 5 showed malformations. The malformations include anomalous pentaradial symmetry, spines in the interambulacral areas, and non-rhomboidal periprocts covered by three supra-anal plates. These findings suggest the need for monitoring external malformations in populations of *A. dufresnii* and other Echinoidea, as they could provide crucial information about the health and status of these populations. Malformations may serve as biomarkers of environmental stress, pollution, or habitat condition changes, which are essential for effective conservation measures.

Key words: *Arbacia dufresnii*, test, malformations, Golfo Nuevo, Echinoidea

Resumen: Malformaciones externas del erizo de mar verde *Arbacia dufresnii* (Blainville, 1825) en poblaciones del Golfo Nuevo, Patagonia, Argentina. Las malformaciones en los erizos de mar pueden manifestarse de diversas formas, y suelen depender de factores ambientales y biológicos a los que estén expuestos. Los adultos de *Arbacia dufresnii* (Blainville, 1825) poseen simetría pentámera, espinas ubicadas en las áreas ambulacrales y la ausencia de estas en las áreas interambulacrales, así como un periprocto de forma romboidal compuesto por cuatro osículos. El objetivo de este reporte es documentar la presencia de malformaciones externas del erizo de mar verde *Arbacia dufresnii* en la población del Golfo Nuevo, Patagonia, Argentina (42°42' S 64°30' O). Los ejemplares fueron recolectados en el Golfo Nuevo. Se recolectaron 319 individuos, 5 de los cuales mostraron malformaciones. Las malformaciones reportadas son: simetría pentámera anormal, presencia de espinas en áreas interambulacrales y periproctos no romboidales cubiertos por tres placas supra-anales. Esperamos que los resultados promuevan el monitoreo de las malformaciones externas en las poblaciones de *A. dufresnii* y otros Echinoidea, lo que podría proporcionar información crucial sobre la salud y estado de las mismas. Las malformaciones podrían ser biomarcadores de estrés ambiental, contaminación o cambios en las condiciones del hábitat, lo cual es fundamental para implementar medidas de conservación efectivas.

Palabras clave: *Arbacia dufresnii*, testa, malformaciones, Golfo Nuevo, Echinoidea

INTRODUCTION

The Class Echinoidea (sea urchins), Holothuroidea (sea cucumbers), Ophiuroidea (brittle, serpent star or ophiuroids), Crinoidea (sea lilies and feathers stars) and Asteroidea (sea stars) constitute the Phylum Echinodermata (echinoderms) (Hickman *et al.*, 2008). This group of animals exhibits numerous distinctive features, including a complex internal transport system known as the water vascular system (WVS), composed of the madreporite plate, which connects the organism's interior to the external environment, and the ambulacral feet, appendages with an ampulla and a podium, with diverse functions (Hickman *et al.*, 2008). Echinoderms possess an endoskeleton composed of numerous calcareous ossicles with accessory structures like spines, spicules, and pedicellariae (Hickman *et al.*, 2008; Borrero Pérez *et al.*, 2012). Sea urchins are capable of regeneration the test and their external appendages (Carnevali, 2006; Emerson *et al.*, 2017; Maboloc & Fang, 2023). They exhibit an indirect life cycle, with embryos and larvae displaying bilateral symmetry (Hickman *et al.*, 2008). Through the process of metamorphosis, involving corporal changes, they acquire both internal and external secondary pentaradial symmetry, although some groups phenotypically display bilateral symmetry (Hickman *et al.*, 2008; Benavides Serrato *et al.*, 2011; Borrero Pérez *et al.*, 2012).

In Echinoidea ossicles are fused to form a test that can exhibit pentaradial symmetry ("regular" sea urchins) or bilateral ("irregular" sea urchins) symmetry (Larrain Prat, 1975; Hickman *et al.*, 2008; Swalla & Smith, 2008; Pérez *et al.*, 2014). The test, consists of twenty columns of ossicles grouped in pairs, forming the five ambulacral areas (AA) and five interambulacral areas (IA), alternating with each other resulting in the juxtaposition of AA with IA (Fig. 1) (Borrero Pérez *et al.*, 2012; Pérez *et al.*, 2014). The AA holds the tube feet of the WVS and spines (Hickman *et al.*, 2008). Each column of calcareous plates extends from the oral area to the apex of the test in the aboral area (Fig. 1) (Borrero Pérez *et al.*, 2012). The apical system is the region where the terminal ossicles of the AA (ocular plate or ambulacra) and IA (genital plate or interambulacral) converge (Fig. 1) (Borrero Pérez *et al.*, 2012; Pérez *et al.*, 2014). Each genital plate has a perforation called genital pore or gonopore through which gametes are released (Fig. 1), each connected to a gonad (Hickman *et al.*, 2008). One of the gen-

ital plates correspond to the madreporite plate of the WVS (Fig. 1) (Hickman *et al.*, 2008; Pérez *et al.*, 2014). The five genital plates are adjacent to the periproct, which is composed of supra-anal calcareous plates (Fig. 1) (Bernasconi, 1925; Hickman *et al.*, 2008). Additionally, sea urchins have a complex chewing called "Aristotle's lantern", composed of five teeth used for cutting and scraping food, which are visible in the oral area (Larrain Prat, 1975; Hickman *et al.*, 2008; Pérez *et al.*, 2014).

Arbacia dufresnii is a sea urchin with a broad distribution range, from the Río de la Plata (Argentina - 35° S) to Puerto Montt (Chile - 42° S), including Antarctica, the Malvinas Islands (Bernasconi, 1925; 1942; Larrain Prat, 1975; Brogger *et al.*, 2013), and smaller islands from the Atlantic Ocean (Caselle *et al.*, 2018). As in other sea urchins, it has ecological significance by influencing productivity in its habitat (Newcombe *et al.*, 2012; Pérez *et al.*, 2014). At the Golfo Nuevo, *A. dufresnii* is currently cultivated in Puerto Madryn by the company ERISEA S.A. for biotechnological purposes. The primary product derived from this species focuses on its female gametes as a source of antioxidants (Rubilar & Cardozo, 2021; Rubilar *et al.*, 2021). During its life cycle, this species typically exhibits primary bilateral symmetry in the embryo/larval development and secondary pentaradial symmetry after metamorphosis (Brogger, 2005; Fernández *et al.*, 2019; Gianguzza, 2020). The test of juveniles and adults are "regular" with well-defined ambulacral and interambulacral areas, and a rhomboidal periproct covered by four ossicles or supra-anal plates (Fig. 1) (Bernasconi, 1925; Larrain Prat, 1975; Brogger, 2005).

Malformations in sea urchins can be manifested in various ways, depending on the environmental and biological factors they are exposed to. These malformations may include alterations in symmetry, such as a reduced or increased number of symmetrical parts, absence or arrangement in spines, anomalies in test plates or in the periproct, and deformities during early development (Hylton Scott, 1939; Larrain Prat, 1975; Munar, 1984; Gambardella *et al.*, 2021). These malformations can be caused by a variety of factors, including chemical pollution, diseases, changes in water temperature and salinity, as well as genetic factors (Dupont *et al.*, 2010; Kroeker *et al.*, 2010; Clark, 2020; Gambardella *et al.*, 2021; Han *et al.*, 2022; González Pisani *et al.*, 2024). The presence of these malformations can serve as an indicator of environmen-




Fig. 1. Diagram of the aboral zone of the test of the sea urchin *Arbacia dufresnii*. AA: ambulacral area; IA: interambulacral area. Detail of the apical zone of the test: genital plates genital pore, ocular plates, and periproct composed of four supra-anal plates. Scale 10 mm.

tal health and habitat quality in marine ecosystems (Laitano *et al.*, 2013; Savriama *et al.*, 2015; Gouveia *et al.*, 2022).

MATERIAL AND METHODS

Sea urchins of *Arbacia dufresnii* were collected during November 2023 by SCUBA diving at Punta Cuevas, Golfo Nuevo, Chubut, Argentina (42°46'44" S, 64°59'52" W); under the regulatory authorization granted by the Ministry of Agriculture and Livestock of the province of Chubut (Disp. N°73/2023–DFyFS MAGIyC). Specimens were transported to the laboratory and acclimated for 15 days in aquaculture recirculating systems supplied with natural seawater in the Experimental Aquarium of the Center for the Study of Marine Systems (CESIMAR CONICET-ARGENTINA), with constant aera-

tion, at 13 ± 0.5 °C and 12:12 light: dark cycle.

All individuals collected were characterized based on their coloration and spines arrangement (Hylton Scott, 1939; Larrain Prat, 1975), and measured for the test height and maximum diameter using a caliper (± 0.1 mm). To determine sex, spawning was induced by injecting 0.3 ml of a 0.5 M KCl saline solution into the peristomial membrane (Brogger, 2005; Fernández *et al.*, 2019). Malformations were identified by comparing the external morphology observed in the organisms test with descriptions from previous authors (Bernasconi, 1925; Larrain Prat, 1975; Brogger, 2005; Epherra, 2010). Specimens displaying any external malformation were drawn and photographed to capture their morphological features in detail, using an iPhone 14 Pro Max and stereoscopic microscope (Stemi 2000, Carl Zeiss) equipped with a digital camera

(Microscopy Service, CCT CONICET-CENPAT).

RESULTS AND DISCUSSION

A total of 319 adults of *Arbacia dufresnii* were collected, 164 females and 150 males in a ratio of 1.09:1; and 5 indetermined specimens. Both test and spine colors characteristic of the species were recorded, according to Larrain Prat (1975). The dominant pigmentation on the test was green (91.11%) with brown spines (85.19%) (Table 1). The test average was 14.53 ± 2.14 mm of height and 28.95 ± 3.04 mm of maximum diameter, which is an intermediate diameter compared to Brogger (2005) at 30.7 ± 4.9 mm and Epherra (2010) at 25.9 mm.

A total of specimens were collected, out of which 5 individuals showed external anomalies in the test, consisting of:

Presence of spines in areas where they should be absent (Bernasconi, 1925; Larrain Prat, 1975):

- Spines in the ossicles' sutures of the interambulacral area (IA) (Fig. 2A, B, C, D). This anomaly was present in 100% of the malformed specimens.
- Spines enclosing the genital plates in the apical area where the IA terminates (Fig. 2B). This was registered found in the 20% of the malformed specimens.

Periproct's malformation (Bernasconi, 1925; Larrain Prat, 1975):

- Periproct was covered by three ossicles (Fig. 2C) instead of four supra-anal plates (Bernasconi, 1925). Larrain Prat (1975) reported this anomaly and detailed that variation of the number of ossicles in the periproct are particular, generally present a higher number of these plates. This anomaly was found in 40% of the malformed specimens.

Symmetry anomaly in the test (Bernasconi, 1925; Larrain Prat, 1975):

- Absence of an ambulacra area (AA) (Fig. 2D). This anomaly was present in 20% of the malformed specimens.

The specimen with anomalous pentaradial symmetry was assessed for maximum diameter using two approaches: AA-IA diameter was measured by placing one caliper point over an ambulacral area and the other over an interambulacral area; IA-IA diameter was measured by position-

ing the caliper between two IA, one positioned in front of the other. The specimen measured 15.22 mm height; the AA-IA diameter was 28.72 mm and the IA-IA diameter was 26.16 mm. On the oral area, the ambulacra and the Aristotle's lantern displayed the typical morphology. At the apex, there were five genital plates, each with a gonopore (Fig. 2D). Upon spawning induction, the organism released gametes through all its gonopores, demonstrating the functionality of its gonads. The released gametes exhibited a whitish pigmentation typical of males to *A. dufresnii* (Brogger, 2005). At the apex, the periproct consisted of four ossicles (Fig. 2D). Unusually, it displayed four AA and four IA (Fig. 2D). Due to the absence of one AA, the IA that would normally be on its sides fused into a single IA that was longer than the others (Fig. 2D). The aboral area showed sparse spines and a vestigial AA (Fig. 2D), suggesting that the AA formation did not complete. Sea urchins are capable of regeneration their external appendages (Carnevali, 2006; Maboloc & Fang, 2023) but the specimen had not sutures on test evidencing regeneration. The regeneration may not produce macroscopic abnormalities (Emerson *et al.*, 2017; Maboloc & Fang, 2023). Consequently, this anomaly positioned the atypical IA in front of two AA and one IA (Fig. 2D), contradicting the typical pattern observed in Echinoidea (Borrero Pérez *et al.*, 2012; Pérez *et al.*, 2014).

This is not the first evidence of absence of an AA in *A. dufresnii* in Argentina. Hylton Scott (1939) reported a test with four AA and four IA, thus naming it tetramorous or tetradial, in Punta Delgada, Patagonia ($42^{\circ}45'59''$ S $63^{\circ}38'13''$ W). However, this specimen reported by Hylton Scott (1939) differs in some aspects from the one described in this study; the AA and IA had a symmetrical length and the test has a cross-like appearance because the AA faces the other AA, and the IA faces other IA. Also, its apical area has four genital plates, one of them with two pores, suggesting it had five gonads. On the other hand, since Hylton Scott (1939) only found the bare test, the sex, number and functionality of the gonads, as well as the symmetry of the oral area, could not be determined. Hylton Scott (1939) suggested that the specimen found at Punta Delgada exhibited internal pentaradial symmetry and external bilateral symmetry. The specimen sampled at Punta Cuevas, reported in this study, exhibits internal pentaradial symmetry. Externally, it showed five teeth in the Aristotle's lantern, five genital plates with




Fig. 2. External malformations found in natural populations of *Arbacia dufresnii* in Golfo Nuevo, Chubut, Argentina (42°46'44" S, 64°59'52" W). A) Specimen with spines in interambulacral areas (IA). B) Specimen with spines along and at the apex of the IA. C) Specimen with spines in IA and periproct covered by three supra-anal plates. D) Specimen with spines in IA and absence of an ambulacral area. Scale 10 mm.

their respective gonopores, and four AA and IA. Therefore, the external symmetry of this specimen cannot be defined as pentaradial or bilateral. These findings could suggest an adverse effect on the metamorphosis process of the specimens, during which pentaradial symmetry is established, although initially the anomaly does not seem to affect either its survival or gamete release.

The presence of spines in the IA has not been reported previously for this species, and there is no literature associating this feature with other Echinoidea. Larrain Prat (1975) described abnormal periprocts composed of three, five or more

ossicles, emphasizing the rarity of this malformation. On the other hand, pentaradial symmetry predominates in Echinodermata, although there are evolutionary asymmetries, such as supernumerary arms in some Asteroidea, external radial symmetry in Holothuroidea or external bilateral symmetry in "irregular" Echinoidea (Hickman *et al.*, 2008; Benavides Serrato *et al.*, 2011; Borrero Pérez *et al.*, 2012). In other cases, external tetraradial symmetry is an atypical morphological feature, as seen in *A. dufresnii* (Echinoidea - Hylton Scott, 1939), *Ophiomyxa pentagona* (Ophiuroidea - Munar, 1984) *Echinaster sepositus*, *Astropecten jonstoni*, *A. aranciacus* and

Table 1. Relative frequency (%) of test colorations in adult sea urchins *Arbacia dufresnii* in the population of Puerto Madryn, Golfo Nuevo, Chubut, Argentina (42°46'44" S, 64°59'52" W).

	Test		
	Pigmentation	Green (%)	Brown (%)
Spines	Brown (%)	76.67	8.52
	Green (%)	11.48	0.37
	Purple (%)	2.96	0.00

Marthasterias glacialis (Asteroidea - Munar, 1984). The limited information on external malformations in Echinodermata complicates comparisons between cases (Hylton Scott, 1939; Munar, 1984; Savriama *et al.*, 2015).

The external malformations of the shell are caused by a variety of factors, such as pollution (Kroeker *et al.*, 2010; Clark, 2020; Han *et al.*, 2022). These malformations can serve as bio-indicators of habitat quality and environmental health in marine ecosystems (Laitano *et al.*, 2013; Savriama *et al.*, 2015; Gouveia *et al.*, 2022). The Mar Argentino (Patagonia, Argentina), the habitat of *Arbacia dufresnii*, is subject to significant anthropic impact caused by tourism, fisheries, and pollution (Brogger *et al.*, 2013; Matteucci & Dadon, 2024). Reported chemical pollutants include heavy metals, microplastics, petrogenic hydrocarbons and others contaminants (Häder *et al.*, 2020; Ríos *et al.*, 2020; Gomez *et al.*, 2024; González Pisani *et al.*, 2024). Despite the low frequency of occurrence of these external malformations (<2%), it is suggested to conduct monitoring to further study them, aiming to detect possible causal factors, analyze their biological effects on organisms and assess their potential as biomarkers of environmental stress.

ACKNOWLEDGEMENTS

This work was supported by the Agencia Nacional de Promoción Científica y Tecnológica (PICT 2018-01729) and Instituto de Conservación de la Ballena (Premio Australis - 2023). We extend special thanks to Téc. Mariano Moris, and Lic. Esteban Nicolás Sarrá for their technical support in the experimental aquarium (CCT CONICET-CENPAT), Téc. Norberto Andrés De Garín for his technical assistance in microscopy (CCT CONICET-CENPAT), to the Nautical Service at CENPAT-CONICET for their collaboration in sampling activities, and especially to “Kevin”, who made this work possible.

REFERENCES

Benavides Serrato, M., G.H. Borrero Pérez & C.M. Diaz Sánchez. 2011. Equinodermos del Caribe colombiano I: Crinoidea, Asteroidea y Ophiuroidea. Instituto de Investigaciones Marinas y Costeras-INVEMAR.

Bernasconi, I. 1925. Resultados de la primera expedición a Tierra del Fuego (1921) enviada por la Facultad de Ciencias exactas, físicas y naturales de la Universidad Nacional de Buenos Aires: Equinodermos. *Anales de la Sociedad Científica Argentina, Equinoideos* (XCIX), 247–257.

Bernasconi, I. 1942. Primeros estadios larvales de *Arbacia dufresnii* (Blv.). *Revista de La Sociedad Argentina de Ciencias Naturales* 19(53): 305–317.

Brogger, M.I. 2005. Biología reproductiva del erizo verde *Arbacia dufresnii* (Blainville, 1825) en costas del Golfo Nuevo, Patagonia. Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires.

Brogger, M.I., D.G. Gil, T. Rubilar, M.I. Martinez, M.E. Díaz de Vivar, M. Escolar, L. Epherra, A.F. Pérez & A. Tablado. 2013. Chapter 11 Echinoderms from Argentina: Biodiversity, Distribution and Current State of Knowledge. In *Echinoderm Research and Diversity in Latin America*, pp. 359–402, Springer-Verlag.

Borrero Pérez, G.H., M. Benavides Serrato & C.M. Diaz Sánchez. 2012. Equinodermos del Caribe colombiano II: Echinoidea y Holothuroidea. Instituto de Investigaciones Marinas y Costeras-INVEMAR.

Carnevali, M.D.C. 2006. Regeneration in Echinoderms: repair, regrowth, cloning. *Invertebrate Survival Journal* 3(1): 64–76.

Caselle, J.E., S.L. Hamilton, K. Davis, C.D.H. Thompson, A. Turchik, R. Jenkinson, D. Simpson & E. Sala. 2018. First quantification of subtidal community structure at Tristan da Cunha Islands in the remote South Atlantic: From kelp forests to the deep sea. *PLoS ONE* 13(3).

Clark, M.S. 2020. Molecular mechanisms of biomineralization in marine invertebrates. *Journal of Experimental Biology* 223.

Dupont, S., O. Ortega Martínez & M. Thorndyke. 2010. Impact of near-future ocean acidification on echinoderms. *Ecotoxicology* 19: 449–462.

Emerson, C.E., H.C. Reinardy, N.R. Bates & A.G. Bodnar. 2017. Ocean acidification impacts spine integrity but not regenerative capacity of spines and tube feet in adult sea urchins. *Royal Society Open Science* 4: 170140.

- Epherra, L. 2010. Ciclo bioquímico del erizo verde *Arbacia dufresnii* (Blainville, 1825), en costas del Golfo Nuevo, Patagonia. Universidad Nacional de la Patagonia San Juan Bosco Facultad de Ciencias Naturales Sede Puerto Madryn.
- Fernández, J.P., L. Epherra, L. Sepúlveda & T. Rubilar. 2019. Desarrollo embrionario y larval del erizo de mar verde *Arbacia dufresnii* (Echinodermata: Echinoidea). *Naturalia Patagónica* 15: 44–58.
- Gambardella, C., F. Marcellini, C. Falugi, S. Varrella & C. Corinaldesi. 2021. Early-stage anomalies in the sea urchin (*Paracentrotus lividus*) as bioindicators of multiple stressors in the marine environment: Overview and future perspectives. *Environmental Pollution* 287: 1–13.
- Gianguzza, P. 2020. Chapter 24: Arbacia. In J.M. Lawrence (Ed.), *Sea urchins: Biology and Ecology*, pp. 419–429, Elsevier.
- Gomez, N.A., J. Sturla Lompré, A. Ferrando, M. Garrido & C.E. Domini. 2024. Update on the status of the contamination by organotin compounds in sediment of Nuevo Gulf, Argentina. Insights from field and experimental studies. *Marine Pollution Bulletin* 200.
- González Pisani, X., J. Sturla Lompré, M. Moris, C. Tropea, L. Stumpf & L. López Greco. 2024. Multiple endpoints analysis of the effects of diesel oil on a commercial species, *Carcinus maenas*. *Science of the Total Environment* 953.
- Gouveia, N., C.A.Y. Harayashiki, F. Márquez, R.A. Lourenço, S. Taniguchi & I.B. Castro. 2022. Mollusc shell shape as pollution biomarkers: Which is the best biological model? *Marine Pollution Bulletin* 179.
- Häder, D.P., A.T. Banaszak, V.E. Villafañe, M.A. Narvarte, R.A. González & E.W. Helbling. 2020. Anthropogenic pollution of aquatic ecosystems: Emerging problems with global implications. *Science of the Total Environment* 713.
- Han, Z., T. Jian, L. Xie & R. Zhang. 2022. Microplastics impact shell and pearl biomineralization of the pearl oyster *Pinctada fucata*. *Environmental Pollution* 293.
- Hickman, C.P., L.S. Roberts, S.L. Keen, A. Larson, H. I'Anson & D.J. Eisenhour. 2008. Chaetognaths, Echinoderms and Hemichordates. In *Integrated Principles of Zoology*, pp. 469–495, McGraw-Hill Higher Education.
- Hylton Scott, M.I. 1939. Un "test" de *Arbacia dufresnii* (Blv) tetrarradiado. *Notas del Museo de La Plata*, 4(16): 1–5.
- Kroeker, K.J., R.L. Kordas, R.N. Crim & G.G. Singh. 2010. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecology Letters* 13: 1419–1434.
- Laitano, M.V., J.D. Nuñez & M. Cledón. 2013. Shell alterations in the limpet *Bostrycapulus odites*: A bioindicator of harbour pollution and mine residuals. *Ecological Indicators* 34: 345–351.
- Larrain Prat, A.P. 1975. Los equinoideos regulares fósiles y recientes de Chile (Vol. 35). *Zoología (Chile)*.
- Maboloc, E.A. & J.K.H. Fang. 2023. Tissue regeneration of the purple sea urchin *Heliocidaris crassispina*. *Bulletin of Marine Science* 99(1): 19–20.
- Matteucci S.D. & J.R. Dadon. 2024. Las zonas costeras y sus recursos naturales: prioridades y perspectivas en Argentina y en el Mundo. En: J.R. Dadon & S.D. Matteucci (ed.), *Zona costera de la pampa argentina: recursos naturales, sostenibilidad, turismo, gestión y derecho ambiental*, pp. 11–30, Nobuko.
- Munar, J. 1984. Anomalías en la simetría de los Asteroidea (Echinodermata). Casos observados en aguas de Mallorca. *Societat d'història Natural de les Balears* 28: 59–66.
- Newcombe, E.M., C.A. Cárdenas & S.W. Geange. 2012. Green sea urchins structure invertebrate and macroalgal communities in the Magellan Strait, southern Chile. *Aquatic Biology* 15: 135–144.
- Pérez, A.F., D.G. Gil & T. Rubilar. 2014. Echinodermata. En J.A. Calcagno (Ed.), *Los Invertebrados Marinos*, pp. 295–316, Fundación de Historia Natural Félix de Azara.
- Ríos, M.F., R.D. Hernández Moresino & D.E. Galván. 2020. Assessing urban microplastic pollution in a benthic habitat of Patagonia Argentina. *Marine Pollution Bulletin* 159.
- Rubilar, T. & D. Cardozo. 2021. Blue Growth: Sea Urchin Sustainable Aquaculture, Innovative Approaches. *Revista de Biología Tropical* 69(S1): 474–486.
- Rubilar, T., E.S. Barbieri, A. Gazquez & M. Avaro. 2021. Sea Urchin Pigments: Echinochrome A and Its Potential Implication in the Cytokine Storm Syndrome. *Marine Drugs* 19(5): 247.
- Savriama, Y., L.C. Stige, S. Gerber, T. Pérez, P. Alibert & B. David. 2015. Impact of sewage pollution on two species of sea urchins in the Mediterranean Sea (Cortiou, France): Radial asymmetry as a bioindicator of stress. *Ecological Indicators* 54: 39–47.
- Swalla, B.J. & A.B. Smith. 2008. Deciphering deuterostome phylogeny: Molecular, morphological and palaeontological perspectives. *Philosophical Transactions of the Royal Society B* 363: 1557–1568.

Doi: 10.22179/REVMACN.27.874

Recibido: 13-VII-2024

Aceptado: 11-IV-2025