

## Diversity of genus *Phromachocrinus* Carpenter, 1879 of the National Invertebrates collection, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”

Renata M. PERTOSS<sup>1</sup>\* & Mariano I. MARTINEZ<sup>1</sup>

<sup>1</sup>Laboratorio de Ecosistemas Marinos, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” (MACN-CONICET), Av. Ángel Gallardo 470 (C1405DJR), Buenos Aires, Argentina.

\*Email: pertossi.renata@macn.gov.ar

**Abstract:** Professor Irene Bernasconi was a pioneer in Antarctic crinoid research in Argentina, with her work on *Promachocrinus kerguelensis* published in 1932. This study review *Promachocrinus* material housed in the collection of the Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” (MACN-In), including specimens collected by Carcelles in 1929. Through an analysis of new samples from Antarctic expeditions and a review of historical material, three species within the genus were identified: *P. kerguelensis*, *P. fragarius*, and *P. joubini*. These studies update the understanding of genus diversity, previously considered a single circum-Antarctic species, and underscore the importance of integrating molecular and morphological methods for identifying cryptic species. This advancement emphasizes the need to reassess historical biodiversity records and highlights the significance of pioneering researchers such as Bernasconi.

**Key words:** Bernasconi, crinoids, Antarctic, lots, cryptic species

**Resumen:** Diversidad del género *Phromachocrinus* Carpenter, 1879 de la Colección Nacional de Invertebrados, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”. La Profesora Irene Bernasconi fue pionera en la investigación de crinoideos antárticos en Argentina, con su trabajo publicado en 1932 sobre *Promachocrinus kerguelensis*. Este estudio se centra en la revisión y actualización del material de *Promachocrinus* de la colección de invertebrados del Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” (MACN-In), que incluye especímenes recolectados por Carcelles en 1929. A través de un análisis de muestras actuales de campañas antárticas y la revisión de material histórico, se identificaron tres especies dentro del género: *P. kerguelensis*, *P. fragarius* y *P. joubini*. Estos estudio actualizan el conocimiento sobre la diversidad del género, previamente considerado como una sola especie circunantártica, y destaca la importancia de la combinación de métodos moleculares y morfológicos en la identificación de especies crípticas. Este avance refuerza la necesidad de reevaluar los registros históricos de biodiversidad y subraya la relevancia de las contribuciones de investigadores pioneros como Bernasconi.

**Palabras clave:** Bernasconi, crinoideo, Antarctic, lotes, especies crípticas

### INTRODUCTION

Professor Irene Bernasconi pioneered crinoid research in Argentina, dedicating her work to the Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” (MACN). Bernasconi (1934) published an article on echinoderms from the Argentine seas, where she mentioned the Antarctic crinoid *Promachocrinus kerguelensis*. However, it was in 1932 that she published her first and only study focused on this crinoid from South Georgia. The material was collected by A. Carcelles in 1929 and deposited in the

MACN Invertebrate collection (MACN-In 18707, MACN-In 18708).

The genus *Promachocrinus* was established by Carpenter (1879), with *P. kerguelensis* as type species, collected during the Challenger expedition (1872–1876). For many years, it was considered a single circum-Antarctic species. However, through molecular and morphological analyses, McLaughlin *et al.* (2023) have revealed a more complex scenario, that corroborates the existence of multiple cryptic species within the genus (Wilson *et al.*, 2007; Hemery *et al.*, 2012; 2013; Eléaume *et al.*, 2014), identifying eight dis-

tinct species. In light of this, the present study provides a review of the genus *Promachocrinus* in the MACN-In, encompassing the material studied by Bernasconi and new material from Antarctic expeditions.

## MATERIAL AND METHODS

We studied samples of *Phromachocrinus* Carpenter, 1879 from the Invertebrate Collection of the Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” (MACN-In) and samples collected during the Antarctic Summer Campaigns (CAV), from 67 to 454 meters depth. The campaigns took place around the Antarctic Peninsula, South Shetland Islands, and South Orkney Islands. Based on diagnostic characters, species descriptions and following the work of McLaughlin *et al.* (2023), the material was identified, and added to the MACN collection. For the specimens already identified as *P. kerguelensis*, we apply the same protocol, in order to revise their identity. A table was provided with each catalog number, coordinates or geographic areas, depth, and specimen identification (Table 1). The location of each specimen was represented on a map (Fig. 1). Maps were created using QGIS v3.8 Zanzibar software.

## RESULTS

### *Promachocrinus kerguelensis* Carpenter, 1879

**Catalog numbers:** MACN-In 18707, MACN-In 18708, MACN-In 27711, MACN-In 35320, MACN-In 44588, MACN-In 44589, MACN-In 44590, MACN-In 44597, MACN-In 44603 (Fig. 1).

**Diagnosis** (after McLaughlin *et al.*, 2023): Radials 10. Arms 20, with dark pigmented spots, 1–2 mm in diameter scattered along arms, sometimes lacking. Centrodorsal rounded to rounded conical. Cirral sockets rounded, extend to ~1 mm from aboral pole. Cirri up to 4 cm long with 40 cirrals. Proximal and distal three cirrals have length half that of width, other cirrals often reasonably uniform, usually only as long as wide, sometimes two or three times the width; penultimate cirral with small opposing spine, terminal claw tends to be hooked.

**Distribution:** Antarctica, recorded in the Antarctic Peninsula, Scotia Arc, Ross Sea, Weddell Sea, and Davis Sea.

**Depth:** 103–926 m

**Remarks:** The specimens MACN-In 18707 and

MACN-In 18708, studied by Irene Bernasconi (1932), were re-examined and their identification was confirmed.

*Promachocrinus fragarius* McLaughlin, Wilson & Rouse, 2023

**Catalog numbers:** MACN-In 26884, MACN-In 35320/1, MACN-In 35321, MACN-In 35322/1, MACN-In 35324, MACN-In 35325/1, MACN-In 44595, MACN-In 44598, MACN-In 44599, MACN-In 44600, MACN-In 445602 (Fig. 1).

**Diagnosis** (after McLaughlin *et al.*, 2023): Radials 10. Arms 20. Centrodorsal shape rounded-conical, slightly increases in width before curving back to rounded tip, strawberry-like. Cirral sockets mostly in columns, taller than wide, covering centrodorsal to aboral pole, occasionally aboral pole tip bare. Pinnules may have pigmentation, purplish, extending to corresponding brachials. Cirri pigmentation paler if present.

**Distribution:** Antarctica, western Antarctic Peninsula, Scotia Arc, Amundsen Sea, Ross Sea, Davis Sea, and Weddell Sea.

**Depth:** 65–1170 m.

### *Promachocrinus joubini* Vaney, 1910

**Catalog numbers:** MACN-In 44593, MACN-In 44594, MACN-In 44596, MACN-In 44601 (Fig. 1).

**Diagnosis** (after McLaughlin *et al.*, 2023): Radials 10. Arms 20. Arms extend from the body nearly horizontally. Centrodorsal 5.4 mm long, 7.4 mm wide; semispherical. Cirral sockets, in three rows, diameter 0.8–1.3 mm; aboral pole lacks cirral sockets. Cirri not present, though drawn for description, proximal segments wider than long, fourth segment as wide as long, and most after that are 1.5 times longer than wide, with distal cirral pointed. Oral disc region 24.7 mm wide. Radial ossicle 1.2 mm long; axil 3.6 mm long. First syzygy occurs at Br3+4, 2.4 mm wide; second syzygy at Br9+10. IBr1+2 protrudes slightly, unlike IIBr1+2.

**Distribution:** Antarctica, western Antarctic Peninsula, Scotia Arc, Amundsen Sea, Ross Sea, Davis Sea, and Weddell Sea.

**Depth:** 110–1157 m.

## DISCUSSION AND CONCLUSION

The revision of *Promachocrinus* specimens from the Invertebrate Collection of the Museo Argentino de Ciencias Naturales “Bernardino




Fig. 1. Map showing the distribution of the lots of *Promachocrinus*: *P. kerguelensis* (orange), *P. fragarius* (purple) and *P. joubii* (blue). Lots of *P. kerguelensis* that worked Bernasconi and published in 1932 (star, MACN-In 18707, MACN-In 18708).

*Rivadavia*" (MACN-In) expands and updates the material deposited for the class. This study builds upon the pioneering work done in Argentina by Professor Irene Bernasconi, who reviewed Antarctic crinoid *Promachocrinus kerguelensis* Carpenter, 1879. Her work was constrained by the limited taxonomic tools available at the time. By revisiting the historical material and incorporating new samples from recent Antarctic expeditions, this research extends our understanding of the genus within the collection. After this study, three species were identified within the

MACN-In collection, including *P. kerguelensis*, *P. fragarius* McLaughlin, Wilson & Rouse, 2023 and *P. joubini* Vaney, 1910, each characterized by unique morphological features and broad bathymetric ranges.

The genus *Promachocrinus* was comprised for one specie (Eléaume *et al.*, 2014; McLaughlin *et al.*, 2023), with two synonyms: *Promachocrinus vanhoeffenianus* Minckert, 1905, and *P. joubini* Vaney, 1910 (Clark, 1915; Mortensen, 1918; John, 1938; Clark & Clark, 1967). Currently, the genus is divided into 8 species that have, 5

radials and 10 arms, such as *P. mawsoni* (Clark, 1937) and *P. wattsonum* McLaughlin, Wilson & Rouse, 2023, or 10 radials and 20 arms, such as *P. kerguelensis*, *P. vanhoeffenianus*, *P. fragarius*, *P. unruhi* McLaughlin, Wilson & Rouse, 2023, *P. uskglass* McLaughlin, Wilson & Rouse, 2023, and *P. joubini*. The genus *Promachocrinus* is endemic to the Southern Ocean, and the specimens analyzed herein, match with the distribution records of McLaughlin *et al.* (2023).

The discovery of these cryptic species necessitates a reevaluation of historical biodiversity records and re-examining collections, using modern analytical tools. What was once perceived as populations of a single species is now recognized as distinct evolutionary lineages (Eléaume *et al.*, 2014). There is evidence of cryptic biodiversity in Antarctica across various taxa (Heimeier *et al.*, 2010; Baird *et al.*, 2011; Brasier *et al.*, 2016; Soler-Membrives *et al.*, 2017; González-Wevar *et al.*, 2019; Jossart *et al.*, 2019; Maroni *et al.*, 2022). This has significant implications for our understanding of Antarctic biodiversity and the evolutionary processes shaping life in these extreme environments. Additionally, the identification of these species raises important questions regarding potential barriers to gene flow and the role of glacial refuge in promoting speciation. The integration of molecular data with traditional morphological approaches has profoundly transformed our understanding of the genus *Promachocrinus*, revealing a previously unrecognized level of diversity. As Antarctic research continues to advance, it is essential to acknowledge the foundational contributions of early researchers, such as Irene Bernasconi, whose work laid the groundwork for these recent discoveries.

#### ACKNOWLEDGMENTS

The authors wish to express their gratitude to Professor I. Bernasconi for her years of dedicated and extensive work on echinoderms. We also thank Gabriela Liuzzi and Mariela Romanelli for providing us with the necessary information for this work. Our thanks extend to the colleagues aboard the B/O Puerto Deseado for their assistance during the Antarctic campaign. Finally, we would like to thank reviewers for valuable suggestions and commentaries that improved this manuscript.

#### REFERENCES

Baird, H.P., K.J. Miller & J.S. Stark. 2011. Evidence of

- hidden biodiversity, ongoing speciation and diverse patterns of genetic structure in giant Antarctic amphipods. *Molecular Ecology* 20: 3439–3454.
- Bernasconi, I. 1932. Notas sobre un crinoideo de Sud Georgia. *Anales del Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”* 37: 29–35.
- Bernasconi, I. 1934. Los equinodermos de los mares argentinos. *Anales del Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”* 53: 1–21.
- Brasier, M. J., H. Wiklund, L. Neal, R. Jeffreys, K. Linse, H. Ruhl & A. G. Glover. 2016. DNA barcoding uncovers cryptic diversity in 50% of deep-sea Antarctic polychaetes. *Royal Society Open Science* 3(11): 160432.
- Carpenter, P. H. 1879. Preliminary report upon the Comatulæ of the Challenger Expedition. *Proceedings of the Royal Society* 28: 383–395.
- Clark, A. H. 1915. Die Crinoiden der Antarktis. *Deutsche-Südpolar Expedition*, 134.
- Clark, A. H. 1937. Crinoidea. *Scientific Reports of the Australian Antarctic Expedition* 8(a): 5–18.
- Clark, A. H. & Clark, A. M. 1967. A monograph of the existing crinoids, Volume 1. The comatulids. Part 5 Suborders Oligophreata (concluded) and Macrophreata. *Bulletin of the United States National Museum* 82: 1–860.
- Eléaume, M., L. G. Hemery, N. Ameziane & M. Roux. 2014. Chapter 10.7. Phylogeographic patterns of the Southern Ocean crinoids. En: C. De Broyer, P. Koubbi, H. Griffiths, B. Raymond, C. D’Udekem d’Acoz, A. Van De Putte, B. Danis, B. David, S. Grant, J. Gutt, C. Held, G. Hosie, F. Huettmann, A. Post & Y. Ropert-Coudert (eds.), *SCAR Marine Biodiversity Information Network, Biogeographic Atlas of the Southern Ocean*, pp. 448–455. The Scientific Committee on Antarctic Research.
- González-Wevar, C. A., K. Gérard, S. Rosenfeld, T. Saucède, J. Naretto, A. Díaz, S. A. Morley, P. Brickle & E. Poulin. 2019. Cryptic speciation in Southern Ocean *Aequioldia eightsi* (Jay, 1839): Mio-Pliocene trans-Drake Passage separation and diversification. *Progress in Oceanography* 174: 44–54.
- Heimeier, D., Lavery, S. & Sewell, M. A. 2010. Molecular species identification of *Astrotruma agassizii* from planktonic embryos: further evidence for a cryptic species complex. *Journal of Heredity* 101(6): 775–779.
- Hemery, L. G., M. Eléaume, V. Roussel, N. Améziane, C. Gallut, D. Steinke, C. Cruaud, A. Couloux & N. G. Wilson. 2012. Comprehensive sampling reveals circumpolarity and sympatry in seven mitochondrial lineages of the Southern Ocean crinoid species *Promachocrinus kerguelensis* (Echinodermata). *Molecular Ecology* 21(10): 2502–2518.
- Hemery, L. G., M. Roux, N. Ameziane & M. Eléaume. 2013. High-resolution crinoid phyletic inter-relationships derived from molecular data. *Marine Biology* 54: 511–523.
- John, D. D. 1938. Crinoidea. *Discovery Reports* 18: 121–222.
- Jossart, Q., C. J. Sands & M. A. Sewell. 2019. Dwarf brooder versus giant broadcaster: combining ge-

Table 1. *Promachocrinus* deposited at MACN-In. (\*) The wet samples were conserved in 96° ethanol, and others in formaldehyde, then in 70° ethanol.

| MACN-In | Family      | Genus                  | Species              | Locality                                  | Depth (m) | Specs. (*) | Conservation          | Id.        | Observation                                                                                                                 |
|---------|-------------|------------------------|----------------------|-------------------------------------------|-----------|------------|-----------------------|------------|-----------------------------------------------------------------------------------------------------------------------------|
| 18707   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | South Georgia                             | 22        | 4          | Dry and wet           | Bernasconi | The lot is divide in a dry sample into the a box (picture of Bernasconi, 1932) and wet samples. Orcadas island cruise, 1978 |
| 18708   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | South Georgia                             | 28.4      | 3          | Dry and wet, EtOH 70° | Bernasconi | Coll. Carcelles y Romero. 1929                                                                                              |
| 27711   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | South Georgia                             |           | 6          | Dry                   |            | Divided into boxes and bags                                                                                                 |
| 35320   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | South Georgia                             |           | 1          | Wet                   |            | Orcadas island cruise, 1978                                                                                                 |
| 44588   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | 62°18' S - 60°02' W                       | 67        | 1          |                       | Pertossi   | E8 L6                                                                                                                       |
| 44589   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | 62°43' S - 56°32' W                       | 202       | 1          |                       | Pertossi   | E22 L28                                                                                                                     |
| 44590   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | 62°10' S - 58°3' W                        | 105       | 2          |                       | Pertossi   | E16 L21                                                                                                                     |
| 44597   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | 62°27,4' S - 61°21,9' W                   | 187       | 1          |                       | Pertossi   | E6 L2                                                                                                                       |
| 44603   | Antedonidae | <i>Phromachocrinus</i> | <i>kerghuelensis</i> | 61°56' S - 58°48' W                       | 206       | 1          |                       | Pertossi   | E11 L11                                                                                                                     |
| 26984   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | Melchior Archipelago, Antarctic Peninsula | 20-50     | 1          | Wet                   | Bernasconi | Coll. Bellisio, 1966                                                                                                        |
| 35320-1 | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | South Georgia                             | 30        | 3          | Wet                   | Pertossi   | Orcadas island cruise, 1978                                                                                                 |
| 35321   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | Zavodovski Island, Scotia Arc             |           | 2          | Wet                   | Pertossi   | Orcadas island cruise, 1978                                                                                                 |
| 44591   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 62°10' S - 58°3' W                        | 105       | 2          | Wet, EtOH 70°         | Pertossi   | E16 L20                                                                                                                     |
| 44592   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 61°54' S - 57°24' W                       | 113       | 1          |                       | Pertossi   | E14 L18                                                                                                                     |
| 44595   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 62°43' S - 56°32' W                       | 202       | 7          |                       | Pertossi   | E22 L28                                                                                                                     |
| 44598   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 62°43' S - 56°30' W                       | 222       | 2          |                       | Pertossi   | E22 L29                                                                                                                     |
| 44599   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 61°51' S - 57°33' W                       | 188       | 3          | Wet, EtOH 70°         | Pertossi   | E13 L15                                                                                                                     |
| 44600   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 61°47' S - 58°26' W                       | 246       | 1          |                       | Pertossi   | E12 L14                                                                                                                     |
| 44602   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | 62°47' S - 60°12' W                       | 454       | 2          | Wet, EtOH 70°         | Pertossi   | E20 L26                                                                                                                     |
| 35325   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | South Shetland Islands                    |           | 1          | Wet                   | Pertossi   | Coll. Cazzaniga. 1976                                                                                                       |
| 35322   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | South Sandwich Islands                    |           | 2          | Wet                   | Pertossi   | Coll. Aldo Toma "Hero", 1973                                                                                                |
| 35324   | Antedonidae | <i>Phromachocrinus</i> | <i>fragarius</i>     | South Shetland Islands                    |           | 3          | Wet                   | Pertossi   | Orcadas island cruise, 1978                                                                                                 |
| 44593   | Antedonidae | <i>Phromachocrinus</i> | <i>joubini</i>       | 62°47' S - 60°12' W                       | 454       | 2          | Wet, EtOH 70°         | Pertossi   | E20 L26                                                                                                                     |
| 44594   | Antedonidae | <i>Phromachocrinus</i> | <i>joubini</i>       | 61°54' S - 57°24' W                       | 113       | 1          |                       | Pertossi   | E14 L18                                                                                                                     |
| 44596   | Antedonidae | <i>Phromachocrinus</i> | <i>joubini</i>       | 62°52,4' S - 60°35,6' W                   | 214       | 1          |                       | Pertossi   | E19 L25                                                                                                                     |
| 44601   | Antedonidae | <i>Phromachocrinus</i> | <i>joubini</i>       | 62°47' S - 60°12' W                       | 454       | 9          | Wet, EtOH 70°         | Pertossi   | E20 L26                                                                                                                     |

- netic and reproductive data to unravel cryptic diversity in an Antarctic brittle star. *Heredity* 123: 622–633.
- Maroni, P. J., B. J. Baker, A. L. Moran, H. A. Woods, C. Avila, G. J. Johnstone, J. S. Stark, K. M. Kocot, S. Lockhart, T. Saucède, G. W. Rouse & N. G. Wilson. 2022. One Antarctic slug to confuse them all: the underestimated diversity of *Doris kerguelensis*. *Invertebrate Systematics* 36(5): 419–435.
- McLaughlin, E. L., N. G. Wilson & G. W. Rouse. 2023. Resolving the taxonomy of the Antarctic feather star species complex *Promachocrinus 'kerguelensis'* (Echinodermata: Crinoidea). *Invertebrate Systematics* 37(7): 498–527.
- Minckert, V. W. 1905. Das Genus *Promachocrinus*, zugleich ein Beitrag zur Faunistik der Antarktis. [The genus *Promachocrinus*, at the same time a contribution to the faunistics of the Antarctic.] *Zoologischer Anzeiger* 28: 490–501.
- Mortensen, T. H. 1918. The Crinoidea of the Swedish Antarctic Expedition. *Wissenschaftliche Ergebnisse der Schwedischen Südpolar-Expedition* 8: 1–23.
- Soler-Membrives, A., K. Linse, K. J. Miller & C. P. Arango. 2017. Genetic signature of last glacial maximum regional refugia in a circum-Antarctic sea spider. *Royal Society Open Science* 4: 170615.
- Vaney, C. 1910. Une nouvelle espèce de *Promachocrinus*. [A new species of *Promachocrinus*.] *Bulletin du Muséum National d'Histoire Naturelle Paris* 16: 158–162. [En français].
- Wilson, N. G., R. L. Hunter, S. J. Lockhart & K. M. Halanych. 2007. Multiple lineages and absence of panmixia in the 'circumpolar' crinoid *Promachocrinus kerguelensis* from the Atlantic sector of Antarctica. *Marine Biology* 152: 895–904.

Doi: 10.22179/REVMACN.27.889

Recibido: 30-IX-2024

Aceptado: 27-I-2025